

ANCESTRY IN LATIN AMERICA: BRAZIL STUDY

Name:

Course:

Institution:

Date:

Abstract

In this systematic review, the researcher sought to investigate the ancestry of populations in Latin America using the case of Brazil. In particular, the review focused on analyzing several pieces of empirical evidence on the genetic history of Brazilians. The study established that the Brazilian population is one of the world's most heterogeneous populations, comprising of Europeans, Amerindians, and Africans. Multiple studies that have employed genetic markers and autosomal polymorphisms, particularly the electrophoresis protein markers and blood groups, indicate that regardless of their skin color, most of the Brazilians exhibit an overwhelming degree of European ancestry, significant scope of African ancestry, and a uniform degree of Amerindian ancestry. The implication of this empirical finding is that the genetic history of the Brazilian people is largely drawn from the European continent.

Keywords: ancestry, Brazil, Europeans, Amerindians, Africans, markers.

ANCESTRY IN LATIN AMERICA: BRAZIL STUDY

Introduction

Genetic association studies (GAS) are usually carried out to establish the specific genetic factors that underlie the vulnerability to complex diseases [1]. Coelho et al. [1] explain that the single nucleotide polymorphisms (SNPs) constitute the major genetic distinctions examined in GAS. Empirical evidence involving GAS and in particular SNP genotyping shows that the Northeastern Brazilian populations experience a significant influence of the European ancestry, followed by the African and Amerindian ancestries respectively [1-3]. Coelho et al.'s [1] study reveals that the dominance of European ancestry on the Northeastern Brazilian populations is 59.7% whereas Campanella et al.'s [2] research indicates that the predominance of the same ancestry in this population is 67.5%. Despite this disparity in the findings, the consensus of both studies is that the

European ancestry has had a considerable impact on the people living in the north east region of Brazil.

It is worth pointing out that other than Coelho et al. [1] and Campanella et al.'s [2] studies, several other pieces of empirical evidence have demonstrated that it is not only the population in the northeastern region of Brazil, but nearly in all of Brazil, whose ancestry is dominated by that q Europeans and Africans [4-6]. Therefore, based on these studies, it can be inferred with ut an prejudice that the majority of the inhabitants in Northeastern Brazil, and Brazil in general, biginated from the European continent or the African continent [4,5]. Nevertheless, this conclus n is not indicative of the assumption that the influence of the Amerindian and Asian and Brazilian population is insignificant. It only implies that the ancestral origin of the najority of e vast Brazilian inhabitants can be traced back to Europe and Africa respectively [6]. othe words, the genetic history of most Brazilians has its roots in both Europe and Africal with the European ancestry showing much greater dominance than the African ancestry. The of this systematic focus review is to explore the ancestry in Latin America using the case study of

Methodology

Search strategy and selection criteria

Typically, a systematic review focuses on identifying, appraising and synthesizing various pieces of evidence that satisfy the eligibility criteria so as to fectively respond to a particular research question. The search strategy that was used in this sy mate review followed the existing guidelines and, more importantly, the researche ulated and adopted a search strategy that is highly detailed, reproducible, and documentable . The esearcher conducted in-depth, objective, and reproducible search of a broad range of sou ses so as to identify as many relevant empirical ms in Nuded "genomic," "ancestry," "Brazil," "Latin pieces of evidence as possible. The sear in the netic," "allelic," "frequencies," "lineages," America," "admixture," population," "mar erg "molecular," "skin color," and "scillen, clusion criteria employed in the search and еi a." post-test data. The search was limited to those articles selection strategy involved precest ar published from 2007 to date. The evalusion criteria limited the search to the full-text articles that were presented in the scientific data ases in English. The PRISMA diagram below (Figure 1) shows the various steps of the march and selection process.

The quality control on the selected articles was carried out by two peer reviewers. They examined the accuracy of the quantitative and qualitative data as well as the research methodologies and the same sizes using a customized Downs and Black checklist.

2)





Multiple genetic studies on normal and pathological distinctions in the human pigmentation have been carried out by many scholars in the recent past [3,7,8]. Some of these pieces of empirical evidence have shown that it is possible to use genotype data to predict the human phenotypes with varying degrees of precision, highlighting the significance of these types of research for forensic practices [3]. Moreover, other studies have suggested that some of the skin pigmentation variations also confer resistance or vulnerability to skin cancer because the presence of different skin tones is mostly attributed to adaptation to diverse environments. It is critical to note that there is a general consensus among human geneticists and anthropologists that from a biological point of view, there are no human races in the world [9-11]. The physical characteristics of a person, in particular, skin pigmentation, shape of nose and lips, hair texture, and hair color are consistently used to group people [10]. Thus, these traits play an extremely important role in human social relationships [9]. Nonetheless, these oligogenic attributes are considered to have developed due to adaptation mechanisms to the geographically dependent environmental factors, for example, solar learned radiation. Therefore, the use of these traits as "race" markers is erroneous [10, 11]. Regardless of this biological meaning, the human skin color remains an important marker that is used to trace ancestry of populations across the world, including Brazil.

Brazil has a relatively large level of genetic admixture [3, 6, 12, 13]. In fact, Br zilians re considered one of the world's most heterogeneous populations, whose ancest com rises of Europeans, Amerindians, and Africans [3]. In a study that sought to establish the color and the genomic ancestry of the Brazilian population, Pimenta et al. [11] observe that based on the estimates of molecular markers, color is a poor predictor of genomic ances. the Brazilian population. The researchers explicate that microsatellites exhibit a high degree of polymorphism because of the variation in the number of the successive units etween the alleles. According to Pimenta et al. [11], this phenomenon is mainly believed to be a alt c recurrent slippage during the replication process. Due to the high mutation rate of the nicrostanites and the step-by-step mutation process results in frequent homoplasy in the population studies. Traditionally, if one is to take into consideration the potential of size constraint and be growth of microsatellites, various populations would tend to depict typical allelic distribution for the markers. Nonetheless, microsatellites have been found to be useful in studying the volution of human beings as well as their genetic structure.

In multiple studies that have used DNA markers to determine the ancestry of Brazilian populations it was found that regardless of their regional curve, we Y chromosomes in the vast majority of the Brazilian males are of European origin [2,9, 1,20]. At the same time, these individuals have an insignificant frequency of the Sup-Salaran version Y chromosomes [2]. Furthermore, the signs of Amerindian ancestry are compately ubsent. The inference that can be drawn from these studies where DNA markers have been used is that there is a strong directional mating that involves the European males and the African and Amerindian females [9]. It is worthwhile pointing out that since Brazil has as a country has a large territory and because several of its diverse population groups moved into different part of the country, it has significant phylogeographical heterogeneity [9].

A critical evaluation of admixture that can be a ena et al.'s [4] empirical observation indicates that Brazil has a significant e appreciated not only at the individual level, but at the group level as well. ack, wledge that at the individual level. Brazilian populations show great ancestry The resear or all the color categories across all the regions [4]. On the other hand, it emerges from variation we khat the African ancestry of the Black individuals is far lower than that witnessed at the group the individual yvel. If one is to take into account the uniqueness of the history and social culture of the Brazilian populations, it becomes clear why there is a significant disparity in both the individual and group admixtures [4]. Africans typically have black skin that is associated with other dominant attributes of color, for example, black curly hair, thicker lips, and broad nose [4,21]. The Europeans have white skin that has a correlation with specific components such as thin nose and lips, lightcolored hair, and light straight hair [4]. According to Pena et al. [4], these characteristics in both Europeans and Africans are all determined by a relatively small number of genes resulting from evolution under certain aspects of the geographical environment, specifically, the levels of ultra violet (UV) exposure. Therefore, if the social identification system is based mainly on the phenotype of individuals, this classification is only based on the presence of specific alleles that are relatively small in number but have an impact on an individual's color, while not taking into account the rest of the genome. On the contrary, if one considers a population that is an outcome of an

extensive admixture of Africans and Europeans, for example, the Brazilian population, the link between color and ancestry should dissipate with time, an observation that has been confirmed by empirical evidence [4,9].

Genomic Ancestry of Brazilian Population

It emerges from a review of the existing body of knowledge on color and genomic ances ty of Brazilian populations that the majority of the researchers have consistently employed the storical. anthropological, and sociological methodology to evaluate the origins of the Brazilia [9]. Contemporary studies that have used uniparental markers in both the black and zilians show that the genomic structure of the majority of the Brazilians is mosaic, con prisin of Nonrecombining Region of the Y Chromosome (NRY) and mtDNA of different hylogeographical origins [9]. Pena et al. [9] point out that the studies with the autosomal big arental markers indicate elevated degrees of genetic admixture between the European, African, a d Amerindian mixtures. However, the researchers observe that it is also clear from these studies to ere was a significant population effect of the "whitening" program that came about during the immigration of the Europeans to Latin America after 1872.

Subsequently, the relationship between genomic ancestry and skin color is imperfect [9, 21]. In other words, it is hard to accurately predict an individual's sk color from their level of Amerindian, African, and European ancestry [9,21]. Therefore, rega ss of beir skin color, most of the Brazilians exhibit an overwhelming degree of Europe n a centry, significant African ancestry, and she, the high ancestral distinction a uniform degree of Amerindian ancestry [9,21]. In a hazian has a singular and fairly individual fraction observed in Blacks and Whites suggests every n their mosaic genomes [9,21]. Understanding the of Amerindian, African, and European ancestre premise of variety in human physical appearance basebeen a subject of long-standing examination ary qualities of physical appearance in human ree intrigue. Refining our knowledge on the b populaces is of impressive development, omedical, and legal significance. Latin America gives a beneficial setting in which to a alyze the test of ancestry on physical appearance [3,8,24]. The locale has a one of a kind hist vider fying with the social and socio-governmental issues of ethnicity, race, and country. An increasive number of hereditary examinations have studied the admixture in Latin America [23,24,2 Barely any studies have been conducted on the effect of hereditary lineage on physical appearance or the relationship of these to singular ideas of ethnicity and family line. Since the tifteenth century, the number of inhabitants in what is currently called "Latin America" has experienced real statistic changes [18,22,24-28]. There have additionally been various discernible population locale of the world 4,19]. ace c anges in the area, Latin America now being the most urbanized

Discussion

The interject of family maps obtained are consistent with other ancestry studies and with evaluation of data on dispersion of fundamental ethnicity bunches inside every nation [5,7,23,27]. This information underlines the strong genetic bonds existing inside and between Latin American nations. Comprehensively, Local American heritage is most noteworthy in territories that were thickly populated in pre-Columbian times and in areas that got generally little non-local migration and which right now have moderately low populace densities [9,13,22]. Early Iberian migrants settled over the mainland, admixing broadly with Local Americans and Africans [21].

Observational confirmation including GAS and specifically SNP genotyping demonstrates that the Northeastern Brazilian populaces have European lineage, followed by the African and Amerindian ancestries individually. One study demonstrates that the frequency of European ancestry in Northeastern Brazilian populaces is 59.7% though another exploration shows that the transcendence of this parentage in this populace is 67.5%. In spite of this dissimilarity, the

European lineage has considerably affected the general population living in the north east district of Brazil.